

# Cheap Nursing CEUs

## Dementia and Sexual Behavior

**1. Dementia is a normal part of aging.**

- A. True
  - B. False
- 

**2. Which of the following statements is most accurate?**

- A. Fronto-temporal dementia is a type of dementia which does not primarily affect the regions of the brain associated with planning, social behavior, and language perception.
  - B. Fronto-temporal dementia is not associated with a younger age of onset, when compared to other types of dementia.
  - C. Vascular dementia may result from strokes and/or other issues that affect blood flow to the brain.
  - D. Vascular dementia is caused by Lewy bodies.
- 

**3. A patient has questions regarding Alzheimer's disease. Which of the following counseling points should be expressed to the patient?**

- A. Alzheimer's disease is not, typically, progressive.
  - B. Alzheimer's disease is, typically, progressive.
  - C. Alzheimer's disease, typically, affects individuals under the age of 60.
  - D. Alzheimer's disease, typically, only affects individuals over the age of 85.
- 

**4. Individuals exhibiting ISB may engage other individuals in conversations regarding sex or make comments related to sex.**

- A. True
  - B. False
- 

**5. Which of the following statements is most accurate?**

- A. Individuals exhibiting dementia associated ISB do not, typically, attempt to touch or grab other individuals.
  - B. Individuals exhibiting dementia associated ISB may attempt to touch or grab other individuals.
  - C. Public disrobing is not considered ISB.
  - D. Public disrobing is only considered to be ISB when it involves individuals over the age of 85.
-

**6. Which of the following recommendations may help health care professionals manage patients exhibiting ISB?**

- A. Health care professionals should indulge patients' "foul language."
  - B. Health care professionals should positively reinforce patients' "foul language."
  - C. Health care professionals should positively reinforce ISB.
  - D. Health care professionals should not positively reinforce ISB.
- 

**7. Due to the nature of dementia, patients may exhibit ISB at specific times of day such as during bathing.**

- A. True
  - B. False
- 

**8. Which of the following statements is most accurate?**

- A. Health care professionals should understand that patients suffering from dementia associated ISB may also be dealing with other health-related conditions or issues.
  - B. Patients suffering from dementia typically do not have any other health-related conditions or issues.
  - C. Health care professionals should avoid considering patient comorbidities.
  - D. Health care professionals should only consider patient comorbidities when caring for patients under the age of 65.
- 

**9. According to the text, health care organizations should ensure patients suffering from dementia receive adequate nutrition and physical activity.**

- A. True
  - B. False
- 

**10. Which of the following statements is most accurate?**

- A. Cognitive behavioral therapy may refer to a form of psychotherapy which focuses on helping individuals solve problems and create positive outcomes by changing unrealistically negative patterns of thought and behavior.
  - B. Cognitive behavioral therapy may refer to a form of psychotherapy which focuses on helping individuals solve problems and create positive outcomes by changing positive patterns of thought and behavior.
  - C. Cognitive behavioral therapy is a form of psychotherapy which may only be used to treat patients under the age of 65.
  - D. Cognitive behavioral therapy is not a form of psychotherapy.
-

**11. Psychoeducation may refer to the process of providing information and education to patients seeking or receiving treatment for mental health-related conditions.**

- A. True
  - B. False
- 

**12. What is the recommended initial adult dose of Namenda?**

- A. 2.5 mg once daily
  - B. 5 mg once daily
  - C. 10 mg once daily
  - D. 15 mg once daily
- 

**13. A fellow health care professional has questions regarding Aricept. Which of the following points of interest should be expressed to the fellow health care professional?**

- A. Aricept is considered to be an antipsychotic agent.
  - B. Aricept is considered to be a selective serotonin reuptake inhibitor.
  - C. The recommended dose of Aricept for individuals with mild to moderate Alzheimer's Disease is 2 mg or 4 mg once daily.
  - D. The recommended dose of Aricept for individuals with mild to moderate Alzheimer's Disease is 5 mg or 10 mg once daily.
- 

**14. A patient has questions regarding Exelon Patch. Which of the following counseling points should be expressed to the patient?**

- A. The recommended initial dose of Exelon Patch is one Exelon Patch (4.6 mg/24 hours) daily.
  - B. The recommended initial dose of Exelon Patch is two Exelon Patches (4.6 mg/24 hours) daily.
  - C. Gastrointestinal adverse effects, including nausea and vomiting, are only common in pediatric patients.
  - D. Gastrointestinal adverse effects, including nausea and vomiting, are not associated with Exelon Patch.
- 

**15. What is the maximum recommended dose of trazodone (Desyrel)?**

- A. 200 mg per day in divided doses
  - B. 250 mg per day in divided doses
  - C. 350 mg per day in divided doses
  - D. 400 mg per day in divided doses
-