

Cheap Nursing CEUs

Fragile X Syndrome

1. Fragile X syndrome is more common in females.

- A. True
 - B. False
-

2. Which of the following may be considered a physical symptom associated with Fragile X syndrome?

- A. Large head
 - B. Large teeth
 - C. Small ears
 - D. Wide face
-

3. Which of the following may be considered a behavioral, social, and emotional symptom associated with Fragile X syndrome?

- A. Hyperactivity
 - B. An inability to learn new skills
 - C. An inability to efficiently or effectively solve problems
 - D. Stuttering
-

4. Which of the following statements is most accurate?

- A. Often individuals show symptoms of Fragile X syndrome by 2 months to 4 months of age.
 - B. Often individuals show symptoms of Fragile X syndrome by 6 months to 10 months of age.
 - C. The physical symptoms associated with Fragile X syndrome typically develop when an individual is about 6 months of age.
 - D. The physical symptoms associated with Fragile X syndrome typically develop when an individual goes through puberty.
-

5. Which of the following may be considered an early sign/symptom of Fragile X syndrome?

- A. Begins to speak early
 - B. Learns to walk early
 - C. Displays unusual reactions to the way things sound, smell, taste, look, or feel
 - D. Enlarged testicles in males
-

6. In order to be diagnosed with an intellectual disability, an individual must have both a significantly low IQ and considerable problems in everyday functioning.

- A. True
 - B. False
-

7. A patient has questions regarding Celexa. Which of the following educational points should be expressed to the patient?

- A. Celexa is a SSRI.
 - B. Celexa is an antipsychotic agent.
 - C. Celexa is not associated with nausea or dry mouth.
 - D. A typical dose of Celexa is 100 mg daily.
-

8. A patient has questions regarding Concerta. Which of the following educational points should be expressed to the patient?

- A. Concerta is a SSRI.
 - B. Concerta is an antipsychotic agent.
 - C. A typical starting dose of Concerta for children and adolescents is 80 mg once daily.
 - D. A typical starting dose of Concerta for children and adolescents is 18 mg once daily.
-

9. What is a typical dose of Adderall XR for individuals ages 6 - 17?

- A. 10 mg once daily
 - B. 50 mg once daily
 - C. 100 mg once daily
 - D. 100 mg twice daily
-

10. A genetic counselor is an individual who specializes in genetics and provides patients with counseling, education, and assistance in making decisions regarding genetic-related health care.

- A. True
 - B. False
-