

# Cheap Nursing CEUs

## Influenza: Prevention and Control

### 1. Which of the following recommendations is most accurate?

- A. Routine annual influenza vaccination of all persons aged  $\geq$ 6 months without contraindications continues to be recommended.
  - B. Routine annual influenza vaccination of all persons aged  $\geq$ 8 months without contraindications continues to be recommended.
  - C. Routine annual influenza vaccination of all persons aged  $\geq$ 9 months without contraindications continues to be recommended.
  - D. Routine annual influenza vaccination of all persons aged  $\geq$ 1 year without contraindications continues to be recommended.
- 

### 2. According to the ACIP General Recommendations on Immunization, which of the following statements is most accurate?

- A. Providers should consider observing all patients for 15 minutes after vaccination to decrease the risk for injury should they experience syncope.
  - B. Providers should consider observing all patients for 30 minutes after vaccination to decrease the risk for injury should they experience syncope.
  - C. Providers should consider observing patients with egg allergies only, for 15 minutes after vaccination to decrease the risk for injury should they experience syncope.
  - D. Providers should consider observing patients with egg allergies only, for 30 minutes after vaccination to decrease the risk for injury should they experience syncope.
- 

### 3. Persons with a history of severe allergic reaction to egg (i.e., any symptom other than hives) should be vaccinated in an inpatient or outpatient medical setting under the supervision of a health care provider who is able to recognize and manage severe allergic conditions.

- A. TRUE
  - B. FALSE
- 

### 4. Which of the following statements is most accurate?

- A. Influenza A and B are the two types of influenza viruses that cause epidemic human disease.
  - B. Influenza A and C are the two types of influenza viruses that cause epidemic human disease.
  - C. Influenza A and D are the two types of influenza viruses that cause epidemic human disease.
  - D. Influenza C and D are the two types of influenza viruses that cause epidemic human disease.
-

**5. Influenza B viruses are separated into two distinct genetic lineages (Yamagata and Victoria) and are categorized into subtypes.**

- A. TRUE
  - B. FALSE
- 

**6. Which of the following statements is most accurate?**

- A. Deaths associated with influenza are most frequent among younger adults.
  - B. Deaths associated with influenza are most frequent among older children.
  - C. Deaths associated with influenza are most frequent among adults below the age of 65.
  - D. Deaths associated with influenza are most frequent among older adults.
- 

**7. Pregnant women are vulnerable to severe symptoms and illness attributable to influenza.**

- A. TRUE
  - B. FALSE
- 

**8. Immune responses elicited by influenza vaccines are generally not strain-specific.**

- A. TRUE
  - B. FALSE
- 

**9. Which of the following statements is most accurate?**

- A. Inactivated influenza vaccines may only be administered by intramuscular injection.
  - B. Inactivated influenza vaccines may only be administered by intradermal injection.
  - C. Inactivated influenza vaccines may be administered by intramuscular or intradermal injection and contain nonreplicating virus.
  - D. Inactivated influenza vaccines may be administered by intramuscular or intradermal injection and do not contain nonreplicating virus.
- 

**10. Vaccine effectiveness is lower among children aged less than 5 years who have never received influenza vaccine previously or who received only 1 dose in their first year of vaccination, than it is among children who received 2 doses in their first year of being vaccinated.**

- A. TRUE
  - B. FALSE
- 

**11. One dose of inactivated influenza vaccine tends to not be immunogenic in healthy adults aged less than 65 years.**

- A. TRUE

B. FALSE

---

**12. According to the text, HIV-infected persons with minimal AIDS-related symptoms and normal or near-normal CD4+ T-lymphocyte cell counts who receive inactivated influenza vaccines develop adequate antibody responses.**

- A. TRUE
  - B. FALSE
- 

**13. Which of the following statements is most accurate?**

- A. Live-attenuated influenza virus strains replicate in nasopharyngeal epithelial cells.
  - B. Live-attenuated influenza virus strains do not replicate in nasopharyngeal epithelial cells.
  - C. Live-attenuated influenza virus strains do not replicate in cells.
  - D. Live-attenuated influenza virus strains do not replicate at all
- 

**14. Which of the following statements is most accurate?**

- A. Fluarix Quadrivalent is indicated in for individuals aged ?3 months.
  - B. Fluarix Quadrivalent is indicated in for individuals aged ?6 months.
  - C. Fluarix Quadrivalent is indicated in for individuals aged ?3 years.
  - D. Fluarix Quadrivalent is indicated in for individuals aged ?8 years.
- 

**15. What is the recommended route of administration for Fluzone Intradermal Quadrivalent?**

- A. IM
  - B. ID
  - C. Subcutaneous
  - D. Oral
- 

**16. Currently available inactivated influenza vaccines are classified as either Pregnancy Category B or Category C.**

- A. TRUE
  - B. FALSE
- 

**17. What is oculorespiratory syndrome?**

- A. An acute, self-limited reaction to inactivated influenza vaccines.
  - B. A chronic, self-limited reaction to inactivated influenza vaccines.
  - C. An acute, self-limited reaction to live-attenuated influenza vaccines.
  - D. A chronic, self-limited reaction to live-attenuated influenza vaccines.
-

**18. Children aged 6 months through 8 years who require 2 doses should receive their first dose as soon as possible after vaccine becomes available, and the second dose ?4 weeks later.**

- A. TRUE
  - B. FALSE
- 

**19. Which of the following statements is most accurate?**

- A. ACIP recommends that all women who are pregnant or who might be pregnant in the upcoming influenza season receive live-attenuated influenza vaccines.
  - B. ACIP recommends that all women who are pregnant or who might be pregnant in the upcoming influenza season receive inactivated influenza vaccines.
  - C. ACIP recommends that women over the age of 40 who are pregnant or who might be pregnant in the upcoming influenza season receive live-attenuated influenza vaccines.
  - D. ACIP recommends that women over the age of 40 who are pregnant or who might be pregnant in the upcoming influenza season receive inactivated influenza vaccines.
- 

**20. Which of the following statements is most accurate?**

- A. Influenza vaccines should be protected from light and stored at recommended temperatures.
  - B. Most influenza vaccines should not be refrigerated.
  - C. Most influenza vaccines should be frozen.
  - D. Multi-dose vials of influenza vaccines should not be returned to recommended storage conditions between uses.
- 

Copyright © 2025 Cheap Nursing CEUs

Visit us at <https://www.cheapnursingceus.com>