

# Cheap Nursing CEUs

## Pediatric Pain Management

### Evaluation of the Supporting Male Survivors of Violence Demonstration Initiative

**1. Pain can refer to an unpleasant sensory and emotional experience associated with actual and potential tissue damage.**

- A. True
  - B. False
- 

### Appendix A: NIJ Research and Evaluation on Children Exposed to Violence-Program-Specific Information

**2. Which of the following statements regarding chronic pain is most accurate?**

- A. Chronic pain typically has an identifiable temporal and causal relationship to injury or disease
  - B. Chronic pain typically resolves after the body heals
  - C. Chronic pain may exist without any clearly identifiable cause
  - D. Chronic pain is not based on duration
- 

### Reducing Medical Errors and Decreasing Risk

**3. Concentrated sucrose solutions may be used as a pain relief measure in preterm and term newborns up to 1 month of age.**

- A. True
  - B. False
- 

### Background

**4. Which of the following non-pharmacological interventions may be used to relieve pediatric patients' pain?**

- A. Distraction techniques

- B. Breast feeding
  - C. Skin-to-skin contact
  - D. All of the above
- 

## **Specifics of CRM in Mental Health Care**

**5. A pediatric patient is initiated on acetaminophen for pain therapy. What is the recommended pediatric maintenance dose of acetaminophen?**

- A. 30 mg/kg PO every 4 hours
  - B. 10 - 15 mg/kg PO every 4 hours
  - C. 25 mg/kg PO every 12 hours
  - D. 6 - 10 mg/kg PO every 6 - 8 hours
- 

**6. Naproxen is contraindicated in pediatric patients.**

- A. True
  - B. False
- 

## **Clinical Risks in Common with Medical Health Care**

**7. A pediatric patient is initiated on morphine for pain therapy. What is the recommended pediatric starting dose of morphine?**

- A. 0.3 mg/kg PO every 3 - 4 hours
  - B. 0.1 mg/kg PO every 3 - 4 hours
  - C. 1 mg PO every 3 - 4 hours
  - D. 1 - 2 mg PO every 6 - 8 hours
- 

## **Discussion**

**8. An adjuvant agent may be used when a pediatric patient is experiencing Stage 2 pain intensity.**

- A. True
  - B. False
-